

## Impact of home care on place of death, access to emergency departments and opioid therapy in 350 terminal cancer patients

Mario Saugo<sup>1</sup>, Michele Pellizzari<sup>1</sup>, Laura Marcon<sup>2</sup>, Pierpaolo Benetollo<sup>3</sup>, Roberto Toffanin<sup>4</sup>, Pietro Gallina<sup>4</sup>, Giovanna Cecchetto<sup>5</sup>, Guido Miccinesi<sup>6</sup>, Stefano Rigon<sup>1</sup>, Maurizio Cancian<sup>7</sup>, and Daniela Sichetti<sup>8</sup>

<sup>1</sup>Servizio Epidemiologico, <sup>2</sup>Servizio Farmaceutico, and <sup>3</sup>Direzione Sanitaria, ULSS 4;

<sup>4</sup>Distretto Sanitario 1 ULSS 16; <sup>5</sup>UO Lungodegenza ULSS 8; <sup>6</sup>CSPO Firenze; <sup>7</sup>MMG Conegliano Veneto;

<sup>8</sup>Laboratorio di Farmacoepidemiologia, Consorzio Mario Negri Sud, Santa Maria Imbaro (Chieti), Italy

Correspondence to: Dr Mario Saugo,  
Servizio Epidemiologico ULSS 4, Via  
Rasa 9, 36016 Thiene (VI), Italy.  
Tel +39-0445-389298;  
fax +39-0445-389245;  
e-mail [mario.saugo@ulss4.veneto.it](mailto:mario.saugo@ulss4.veneto.it)

Received March 28, 2007;  
accepted September 7, 2007.